

Taq DNA Polymerase 2x Master Mix RED

2 mM MgCl₂ final concentration

MADE IN DENMARK

Cat. No.: A190303

500 Reactions



A190303

-	Taq DNA Polymerase 2x Master Mix RED, 2 mM MgCl ₂
ID No.	5200350
Cap colour	Purple
Content	10 x 1.25 ml

Key Features

Taq DNA Polymerase 2x Master Mix RED is a ready-to-use 2x reaction mix with the Ampliqon Taq DNA polymerase, the NH₄⁺ buffer system, dNTPs and magnesium chloride present. Each reaction requires 25 µl of the 2x Master Mix RED. Simply add primers, template and water to a total reaction volume of 50 µl to successfully carry out primer extensions and other molecular biology applications.

Taq DNA Polymerase 2x Master Mix RED offers several advantages. Set up time is significantly reduced. The chance of contaminating component stocks is eliminated. Reduction of reagent handling steps leads to better reproducibility. Standard tests can be set up with the confidence that results will be consistent every time.

There is no need to buy and use separate loading dyes. Simply load a portion of the reaction product onto an agarose gel for electrophoresis and subsequent visualization. The red dye front runs at 300 – 1000 bp on a 0.5 – 1.5% agarose gel.

Composition of the Taq DNA Polymerase 2x Master Mix RED (2 mM MgCl₂ final concentration)

- Tris-HCl pH 8.5, (NH₄)₂SO₄, 4 mM MgCl₂, 0.2% Tween® 20
- 0.4 mM of each dNTP
- Ampliqon Taq DNA polymerase
- Inert red dye and stabilizer

Recommended Storage and Stability

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label.

Option: Store at +4 °C for up to 6 months.

Quality Control

Taq DNA Polymerase is tested for contaminating activities, with no traces of endonuclease activity, nicking activity or exonuclease activity.

Protocol

This protocol serves as a guideline to ensure optimal PCR results when using Taq DNA Polymerase 2x Master Mix RED. Optimal reaction conditions such as incubation times, temperatures, and amount of template DNA may vary and must be determined individually.

1. Thaw Taq 2x Master Mix RED and primers. **It is important to thaw the solutions completely and mix thoroughly before use to avoid localized concentrations of salts.** Keep all components on ice.
2. Prepare a reaction mix. Table 1 shows the reaction set up for a final volume of 50 µL. If desired, the reaction size may be scaled down. Use 10 µl of the Taq 2x Master Mix RED in a final volume of 20 µl.

Table 1. Reaction components (reaction mix and template DNA)

Component	Vol./reaction*	Final concentration*
Taq 2x Master Mix	25 µl	1x
25 mM MgCl ₂	0 µl (0 – 5 µl)	2 mM (2 – 4.5 mM)
Primer A (10 µM)	1 µl (0.5 – 5 µl)	0.2 µM (0.1 – 1.0 µM)
Primer B (10 µM)	1 µl (0.5 – 5 µl)	0.2 µM (0.1 – 1.0 µM)
PCR-grade H ₂ O	X µl	-
Template DNA	X µl	genomic DNA: 50 ng (10 – 500 ng) plasmid DNA: 0.5 ng (0.1 – 1 ng) bacterial DNA: 5 ng (1 – 10 ng)
TOTAL volume	50 µl	-

* Suggested starting conditions; theoretically used conditions in brackets

3. Mix the reaction mix thoroughly and dispense appropriate volumes into reaction tubes. Mix gently, e.g. by pipetting the reaction mix up and down a few times.
4. Add template DNA to the individual tubes containing the reaction mix.
5. Program the thermal cycler according to the manufacturer's instructions. See table 2 for an example.
For maximum yield and specificity, temperatures and cycling times should be optimized for each new template target or primer pair.
6. Place the tubes in the thermal cycler and start the reaction.
7. At the end of the run, simply load a portion of the reaction product (e.g. 10 µl) onto an agarose gel for analysis.

Table 2. Three-step PCR program

Cycles	Duration of cycle	Temperature
1	2 – 5 minutes	95 °C
25 - 35	20 – 30 seconds ^a 20 – 40 seconds ^b 30 seconds ^c	95 °C 50 – 65 °C 72 °C
1	5 minutes ^d	72 °C

^a. Denaturation step: This step is the first regular cycling event and consists of heating the reaction to 95 °C for 20 – 30 seconds. It causes melting of the DNA template by disrupting the hydrogen bonds between complementary bases, yielding single-stranded DNA molecules.

^b. Annealing step: The reaction temperature is lowered to 50 – 65 °C for 20 – 40 seconds allowing annealing of the primers to the single-stranded DNA template. Typically, the annealing temperature is about 3 – 5 °C below the T_m (melting temperature) of the primers used.

- ^c Extension/elongation step: Taq polymerase has its optimum activity temperature at 72 °C. At this step the DNA polymerase synthesizes a new DNA strand complementary to the DNA template strand. The extension time depends on the length of the DNA fragment to be amplified. As a rule of thumb, at its optimum temperature the DNA polymerase will polymerize a thousand bases per minute.
- ^d Final elongation: This single step is occasionally performed at a temperature of 72 °C for 5 minutes after the last PCR cycle to ensure that any remaining single-stranded DNA is fully extended.

Two-step PCR program

Fast 2-step PCR protocols are available using this link:

<https://ampliqon.com/en/pcr-technology/application-notes/>

Notes:

- The final MgCl₂ concentration of this 2x Taq Master Mix RED is 2 mM. In some applications, more than 2 mM MgCl₂ is required for best results. Use 25 mM MgCl₂ to adjust the Mg²⁺ concentration according to table 3.

Table 3. Additional volume (μl) of MgCl₂ per 50 μl reaction:

Final MgCl ₂ conc. in reaction (mM)	2.0	2.5	3.0	3.5	4.0	4.5
Volume of 25 mM MgCl ₂	0	1	2	3	4	5

Related Products

Taq Master Mixes (500 x 50 μl reactions) *	Cat. No.
2x Master Mix, 1.5 mM MgCl ₂ final concentration	A140303
2x Taq OptiMix CLEAR, 1.5 mM MgCl ₂ final concentration	A370503
2x Master Mix RED, 1.5 mM MgCl ₂ final concentration	A180303

TEMPase Hot Start Master Mixes (500 x 50 μl reactions) *	Cat. No.
2x Master Mix A**, 1.5 mM MgCl ₂ final concentration	A230303
2x Master Mix A**BLUE, 1.5 mM MgCl ₂ final concentration	A290403

*Master mixes available also in 1.1x variants as well as 2 mM MgCl₂ variants, **Mix A is Ammonium Buffer based, also available as Mix C based on Combination Buffer.

Special TEMPase Master Mixes (500 x 50 μl reactions)	Cat. No.
Multiplex 2x Master Mix, 3 mM MgCl ₂ final concentration	A260303
GC TEMPase 2x Master Mix I – for GC-rich templates	A331703
GC TEMPase 2x Master Mix II – for GC-rich templates	A332703

Taq DNA Polymerase (500 units) *	Cat. No.
Taq DNA Polymerase 5 U/μl	A110003
• with 10x Ammonium Buffer	A111103

*Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM MgCl₂

Hot Start DNA Polymerase (500 units) *	Cat. No.
TEMPase Hot Start DNA Polymerase, 5 U/μl	A220003
• with 10x Ammonium Buffer	A221103

*Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM MgCl₂

Buffers for DNA polymerases *	Cat. No.
10x Ammonium Buffer, 3 x 1.5 ml	A301103
10x Standard Buffer, 3 x 1.5 ml	A302103
10x Combination Buffer, 3 x 1.5 ml	A303103
5x PCR Buffer RED, 6 x 1.5 ml **	A301810
PCR Grade Water, 6 x 5 ml	A360056

*Ammonium Buffer, Standard Buffer and Combination Buffer are also available as Mg²⁺ free buffers, detergent free buffers and Mg²⁺ and detergent free buffers.

**For direct gel loading and visualisation.

For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at www.ampliqon.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.

Made in Denmark

Issued 02/2025