

Taq DNA Polymerase RED

Concentration: 5 units/ μ l

MADE IN DENMARK

Cat. No.	Units	Taq DNA Polymerase RED ID: 5101750	10x Ammonium Buffer, 15 mM MgCl ₂ ID: 5100950	MgCl ₂ 25 mM ID: 5575801
A201103	500	100 μ l	1.5 ml	1.5 ml
A201104	1000	2 x 100 μ l	2 x 1.5 ml	2 x 1.5 ml
A201106	2500	5 x 100 μ l	5 x 1.5 ml	5 x 1.5 ml
A201107	5000	10 x 100 μ l	3 x 5 ml	3 x 5 ml
A201108	10,000	3 x 667 μ l	6 x 5 ml	6 x 5 ml

Key Features

Ampliqon Taq DNA Polymerase is a thermostable, recombinant DNA polymerase, which exhibits high activity in primer extension and other molecular biology applications. The enzyme is isolated from *Thermus aquaticus* and has a molecular weight of approximately 94 kDa. Ampliqon Taq DNA Polymerase has a 5'→3' DNA polymerase and a 5'→3' exonuclease activity. Taq DNA Polymerase RED contains a red dye which provides easy and quick identification of reactions to which enzyme was added and allows confirmation of complete mixing. The inert dye has no effect on downstream processes. Taq DNA Polymerase RED is added directly to the reaction mix and is used in the same manner as standard Taq DNA Polymerase.

Kit Components

Ampliqon Taq DNA Polymerase RED in Storage Buffer

5 U/ μ l Taq, 20 mM Tris-HCl pH 8.3, 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, inert dye, 0.5% Tween® 20, 50% glycerol.

10x Ammonium Buffer

Tris-HCl pH 8.5, (NH₄)₂SO₄, 15 mM MgCl₂, 1% Tween® 20.

Ammonium in the buffer minimizes the need for optimization of the MgCl₂ concentration or the annealing temperature for most primer-template systems.

25 mM MgCl₂

Recommended Storage and Stability

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label.

Option: Store at +4 °C for up to 6 months.

Quality Control

Taq DNA Polymerase is tested for contaminating activities, with no traces of endonuclease activity, nicking activity or exonuclease activity.

Unit Definition

One unit is defined as the amount of polymerase that incorporates 10 nmol of dNTPs into acid-precipitable DNA in 30 minutes at 72 °C under standard assay conditions.

Protocol

This protocol serves as a guideline to ensure optimal PCR results when using Taq DNA Polymerase. Optimal reaction conditions such as incubation times, temperatures and amount of template DNA may vary and must be determined individually.

1. Thaw 10x Buffer, dNTP mix and primer solutions. **It is important to thaw the solutions completely (some buffers need to reach room temperature) and mix thoroughly before use to avoid localized concentrations of salts.** Keep all components on ice. The polymerase is provided in glycerol and does not need thawing. Keep it at -20 °C at all times.
2. Set up reaction mixtures in an area separate from that used for DNA preparation or product analysis. Work on ice at all times.
3. Prepare a master mix according to Table 1. The master mix typically contains all the components needed for extension except the template DNA.

Table 1. Reaction components (master mix and template DNA)

Component	Vol./reaction*	Final concentration*
10x Buffer	5 μ l	1x
25 mM MgCl ₂	0 μ l (0-6 μ l)	1.5 mM (1.5-4.5 mM)
dNTP mix (10 mM each)	1 μ l	0.2 mM of each dNTP
Primer A (10 μ M)	1 μ l (0.5-5 μ l)	0.2 μ M (0.1-1.0 μ M)
Primer B (10 μ M)	1 μ l (0.5-5 μ l)	0.2 μ M (0.1-1.0 μ M)
Taq DNA Pol.	0.2 μ l (0.2-1 μ l)	1 unit (1-5 units)
PCR-grade H ₂ O	X μ l	-
Template DNA	X μ l	genomic DNA: 50 ng (10-500 ng) plasmid DNA: 0.5 ng (0.1-1 ng) bacterial DNA: 5 ng (1-10 ng)
TOTAL volume	50 μ l	-

* Suggested starting conditions; theoretically used conditions in brackets. The final volume can be reduced to 25 μ l by using half of the volumes suggested in Vol./reaction, eg. 0.1 μ l Taq instead of 0.2 μ l Taq.

4. Mix the master mix thoroughly and dispense appropriate volumes into reaction tubes. Mix gently, e.g. by pipetting the master mix up and down a few times.
5. Add template DNA to the individual tubes containing the master mix.
6. Program the thermal cycler according to the manufacturer's instructions.
For maximum yield and specificity, temperatures and cycling times should be optimized for each new template target or primer pair.
7. Place the tubes in the thermal cycler and start the reaction.

Three-step PCR program

Cycles	Duration of cycle	Temperature
1	2-5 minutes ^a	95 °C
25-35	20-30 seconds ^b 20-40 seconds ^c 30 seconds ^d	95 °C 50-65 °C 72 °C
1	5 minutes ^e	72 °C

^a. Initial denaturation step (optional).

^b. Denaturation step: This step is the first regular cycling event and consists of heating the reaction to 95 °C for 20-30 seconds. It causes melting of the DNA template by disrupting the hydrogen bonds between complementary bases, yielding single-stranded DNA molecules.

^c. Annealing step: The reaction temperature is lowered to 50-65 °C for 20-40 seconds allowing annealing of the primers to the single-stranded DNA template. Typically, the annealing temperature is about 3-5 °C below the T_m (melting temperature) of the primers used.

^d. Extension/elongation step: Taq polymerase has its optimum activity temperature at 72 °C. At this step the DNA polymerase synthesizes a new DNA strand complementary to the DNA template strand. The extension time depends on the length of the DNA fragment to be amplified. As a rule of thumb, at its optimum temperature the DNA polymerase will polymerize a thousand bases per minute.

^e. Final elongation: This single step is occasionally performed at a temperature of 72 °C for 5 minutes after the last PCR cycle to ensure that any remaining single-stranded DNA is fully extended.

Two-step PCR program

Fast 2-step PCR protocols are available using this link:

<https://ampliqon.com/en/pcr-technology/application-notes/>

Notes:

- 15 mM MgCl₂ is present in 10x PCR Buffer. The 1x concentration is 1.5 mM MgCl₂. In some applications, more than 1.5 mM MgCl₂ is required for best results. For this reason, 25 mM MgCl₂ is included in the kit. Table 2 provides the volume of 25 mM MgCl₂ to be added to the master mix if a higher MgCl₂ concentration is required.

Table 2. Additional volume (μl) of MgCl₂ per 50 μl reaction

Final MgCl ₂ conc. in reaction (mM)	1.5	2.0	2.5		3.0	3.5	4.0	4.5
Volume of 25 mM MgCl ₂	0	1	2		3	4	5	6

- For longer DNA targets: More Taq DNA polymerase could be added to the PCR master mix.

For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at www.ampliqon.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.

Made in Denmark

Issued 06/2025