



## TEMPase Hot Start DNA Polymerase, Glycerol Free

With 10x Ammonium Buffer (15 mM MgCl<sub>2</sub>)

Concentration: 5 units/μl



A241104

Cat. No.: A241104

1000 Units

|            | TEMPase, Glycerol Free,<br>5 U/μl | 10x Ammonium<br>Buffer,<br>15 mM MgCl <sub>2</sub> | MgCl <sub>2</sub><br>25 mM |
|------------|-----------------------------------|--|----------------------------|
| ID No.     | 5101700                           | 5100950  | 5575801                    |
| Cap colour | White                             | White  | Clear                      |
| Content    | 2 x 100 μl                        | 2 x 1.5 ml   | 2 x 1.5 ml                 |

### Key Features

TEMPase Hot Start DNA Polymerase, Glycerol Free, is a high-quality DNA polymerase developed for automation. It is a glycerol free formulation of TEMPase Hot Start Polymerase with the same excellent performance. The glycerol free formulation makes it well suited for automation and freeze-drying.

TEMPase Hot Start DNA Polymerase is a modified form of Ampliqon Taq DNA Polymerase. A chemical moiety is attached to the enzyme at the active site, which renders the enzyme inactive at room temperature. Thus, during setup and the first ramp of thermal cycling, the enzyme is not active and misprimed primers are not extended. Once the reaction reaches optimal activating temperature, the chemical moiety is cleaved during a 15 minutes heat activation step, releasing the active TEMPase Hot Start DNA Polymerase into the reaction.

### Kit Components

#### TEMPase Hot Start DNA Polymerase in Glycerol Free Storage Buffer

5 U/μl TEMPase Hot Start DNA Polymerase, 20 mM Tris-HCl pH 8.9, 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5% Tween® 20.

#### 10x Ammonium Buffer

Tris-HCl pH 8.5, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 15 mM MgCl<sub>2</sub>, 1% Tween® 20.

Ammonium Buffer gives a superior amplification signal (high yield) in most primer-template systems. Ammonium in the buffer minimizes the need for optimization of the MgCl<sub>2</sub> concentration and the annealing temperature.

#### MgCl<sub>2</sub>

25 mM MgCl<sub>2</sub> in PCR grade water.

#### Recommended Storage and Stability

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label.

Option: Store at +4 °C for up to 6 months.

#### Quality Control

TEMPase Hot Start DNA Polymerase is tested for contaminating activities, with no traces of endonuclease activity, nicking activity or exonuclease activity.

### Unit Definition

One unit is defined as the amount of polymerase that incorporates 10 nmol of dNTPs into acid-precipitable DNA in 30 minutes at 72 °C under standard assay conditions.

### Protocol

This protocol serves as a guideline to ensure optimal PCR results when using TEMPase Hot Start DNA Polymerase. Optimal reaction conditions such as incubation times, temperatures and amount of template DNA may vary and must be determined individually.

1. Thaw DNA polymerase, 10x Buffer, dNTP mix and primer solutions. **It is important to thaw the solutions completely (some buffers need to reach room temperature) and mix thoroughly before use to avoid localized concentrations of salts.**

**Important:** Spin vials briefly before use.

2. Prepare a reaction mix according to table 1. The reaction mix typically contains all the components needed for primer extension except the template DNA.

**Table 1. Reaction mix and template DNA**

| Component                  | Vol./reaction*         | Final concentration*   |
|----------------------------|------------------------|--|
| 10x Buffer                 | 5 μl                   | 1x   |
| 25 mM MgCl <sub>2</sub>    | 0 μl (0 – 6 μl)        | 1.5 mM (1.5 – 4.5 mM)  |
| dNTP mix<br>(10 mM each)   | 1 μl                   | 0.2 mM of each dNTP  |
| Primer A (10 μM)           | 1 μl (0.5 – 5 μl)      | 0.2 μM (0.1 – 1.0 μM)  |
| Primer B (10 μM)           | 1 μl (0.5 – 5 μl)      | 0.2 μM (0.1 – 1.0 μM)  |
| TEMPase DNA<br>Pol.        | 0.4 μl<br>(0.2 – 1 μl) | 2 units (1 – 5 units)  |
| PCR-grade H <sub>2</sub> O | X μl                   | -  |
| Template DNA               | X μl                   | genomic DNA: 20 ng (1 – 200 ng)<br>plasmid DNA: 0.5 ng (0.1 – 1 ng)<br>bacterial DNA: 5 ng (1 – 10 ng) |
| <b>TOTAL volume</b>        | 50 μl                  | -  |

\* Suggested starting conditions; theoretically used conditions in brackets. The final volume can be reduced to 25 μl by using half of the volumes suggested in Vol./reaction, eg. 0.2 μl TEMPase instead of 0.4 μl TEMPase.

3. Mix the reaction mix thoroughly and dispense appropriate volumes into reaction tubes.
4. Add template DNA to the individual tubes containing the reaction mix.
5. Program the thermal cycler according to the manufacturer's instructions. **Each program must start with an initial heat activation step at 95°C for 15 minutes.**  
For maximum yield and specificity, temperatures and cycling times should be optimized for each new template or primer pair.
6. Place the tubes in the thermal cycler and start the reaction.

**Table 2. Three-step PCR program**

| Cycles  | Duration of cycle  | Temperature                  |
|---------|--|------------------------------|
| 1       | 15 minutes <sup>a</sup>  | 95 °C                        |
| 25 – 35 | 20 – 30 seconds <sup>b</sup><br>20 – 40 seconds <sup>c</sup><br>30 – 90 seconds <sup>d</sup> | 95 °C<br>50 – 65 °C<br>72 °C |
| 1       | 5 minutes <sup>e</sup>   | 72 °C                        |

<sup>a</sup>. For activation of the TEMPase hot start enzyme.

<sup>b</sup>. Denaturation step: This step is the first regular cycling event and consists of heating the reaction to 95 °C for 20 – 30 seconds. It causes melting of the DNA template by disrupting the hydrogen bonds

between complementary bases, yielding single-stranded DNA molecules.

- c. Annealing step: The reaction temperature is lowered to 50 – 65 °C for 20 – 40 seconds allowing annealing of the primers to the single-stranded DNA template. Typically, the annealing temperature is about 3 – 5 °C below the  $T_m$  (melting temperature) of the primers used.
- d. Extension/elongation step: TEMPase polymerase has its optimum activity temperature at 72 °C. At this step the DNA polymerase synthesizes a new DNA strand complementary to the DNA template strand. The extension time depends on the length of the DNA fragment to be amplified. As a rule of thumb, at its optimum temperature the DNA polymerase will polymerize a thousand bases per minute.
- e. Final elongation: This single step is occasionally performed at a temperature of 72 °C for 5 minutes after the last PCR cycle to ensure that any remaining single-stranded DNA is fully extended.

#### Notes:

- 15 mM  $MgCl_2$  is present in 10x PCR Buffer. The 1x concentration is 1.5 mM  $MgCl_2$ . In some applications, more than 1.5 mM  $MgCl_2$  is required for best results. For this reason, 25 mM  $MgCl_2$  is included in the kit. Table 3 provides the volume of 25 mM  $MgCl_2$  to be added to the master mix if a higher  $MgCl_2$  concentration is required.

**Table 3. Additional volume (µl) of  $MgCl_2$  per 50 µl reaction**

| Final $MgCl_2$ conc. in reaction (mM) | 1.5 | 2.0 | 2.5 | 3.0 | 3.5 | 4.0 | 4.5 |
|---------------------------------------|-----|-----|-----|-----|-----|-----|-----|
| Volume of 25 mM $MgCl_2$              | 0   | 1   | 2   | 3   | 4   | 5   | 6   |

- For longer DNA targets more DNA polymerase could be added to the PCR master mix.

Tween® 20 is a registered trademark of ICI Americas Inc.

## Related Products

| TEMPase DNA Polymerase (500 units) *                   | Cat. No.       |
|--|----------------|
| TEMPase Hot Start DNA Polymerase 5 U/µl                | <b>A220003</b> |
| • with 10x Ammonium Buffer                             | <b>A221103</b> |
| • with 5x PCR Buffer RED                               | <b>A221803</b> |
| TEMPase Hot Start DNA Polymerase 5 U/µl, glycerol free | <b>A240003</b> |
| • with 10x Ammonium Buffer                             | <b>A241103</b> |

| Taq DNA Polymerase (500 units) * | Cat. No.       |
|----------------------------------|----------------|
| Taq DNA Polymerase, 5 U/µl       | <b>A110003</b> |
| • with 10x Ammonium Buffer       | <b>A111103</b> |

\*Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM  $MgCl_2$ .

| Buffers for DNA polymerases *      | Cat. No.       |
|------------------------------------|----------------|
| 10x Ammonium Buffer, 3 x 1.5 ml    | <b>A301103</b> |
| 10x Standard Buffer, 3 x 1.5 ml    | <b>A302103</b> |
| 10x Combination Buffer, 3 x 1.5 ml | <b>A303103</b> |
| 5x PCR Buffer RED, 6 x 1.5 ml **   | <b>A301810</b> |

\*Ammonium Buffer, Standard Buffer and Combination Buffer are also available as  $Mg^{2+}$  free buffers, detergent free buffers and  $Mg^{2+}$  and detergent free buffers.

\*\*For direct gel loading and visualisation.

| TEMPase Hot Start Master Mixes (500 x 50 µl reactions) *   | Cat. No.       |
|--|----------------|
| 2x Master Mix A**, 1.5 mM $MgCl_2$ final concentration     | <b>A230303</b> |
| 2x Master Mix A**BLUE, 1.5 mM $MgCl_2$ final concentration | <b>A290403</b> |

\*Master mixes available also in 1.1x variants as well as 2 mM  $MgCl_2$  variants, \*\*Mix A is Ammonium Buffer based, also available as Mix C based on Combination Buffer.

| Special Master Mixes (500 x 50 µl reactions)               | Cat. No.       |
|--|----------------|
| Multiplex 2x Master Mix, 3 mM $MgCl_2$ final concentration | <b>A260303</b> |
| GC TEMPase 2x Master Mix I – for GC-rich templates         | <b>A331703</b> |
| GC TEMPase 2x Master Mix II – for GC-rich templates        | <b>A332703</b> |

| Ultrapure dNTPs*  | Cat. No.       |
|---|----------------|
| dNTP Mix 40 mM (2 x 500 µl): 10 mM each dA, dC, dG, dT  | <b>A502004</b> |
| dNTP Set, 100 mM each: 250 µl of each dA, dC, dG and dT | <b>A511104</b> |

\*Other concentrations and Single dNTPs are available.

| Loading Buffers, PCR water and Ladders        | Cat. No.       |
|---|----------------|
| 5x Loading Buffer Red *, 5 x 1 ml             | <b>A608104</b> |
| Iqon PCR Ladder **, 100 – 3000 bp, 1 x 0.5 ml | <b>A610341</b> |
| PCR Grade Water, 6 x 5 ml                     | <b>A360056</b> |

\*Also available with Blue, Orange or Cyan. \*\* Available in different size ranges.

For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at [www.ampliqon.com](http://www.ampliqon.com) or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.

**Made in Denmark**

Issued 08/2021