

Taq DNA Polymerase, Glycerol free

With 10x Ammonium Buffer (15 mM MgCl₂)

Concentration: 5 units/μl



A101106

Cat. No.: A101106

2500 Units

	Taq DNA Polymerase Glycerol free 5 U/μl	10x Ammonium Buffer, 15 mM MgCl ₂	MgCl ₂ 25 mM
ID No.	5101500	5100950	5575801
Cap colour	Black	White	Clear
Content	5 x 100 μl	5 x 1.5 ml	5 x 1.5 ml

MADE IN DENMARK

Key Features

Ampliqon Taq DNA Polymerase is a thermostable, recombinant DNA polymerase, which exhibits very high activity in primer extension and other molecular biology applications. The enzyme is isolated from *Thermus aquaticus* and has a molecular weight of approximately 94 kDa. Ampliqon Taq DNA Polymerase has a 5'→3' DNA polymerase and a 5'→3' exonuclease activity. The enzyme lacks a 3'→5' exonuclease activity (no proofreading ability). Taq DNA Polymerase leaves an A' overhang, which makes the enzyme ideal for TA cloning.

Taq DNA Polymerase Glycerol free is ideal for freeze drying and automation.

Kit Components

Ampliqon Taq DNA Polymerase in Storage Buffer, Glycerol free
5 U/μl Taq, 20 mM Tris-HCl pH 8.3, 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.5% Tween® 20.

10x Ammonium Buffer

Tris-HCl pH 8.5, (NH₄)₂SO₄, 15 mM MgCl₂, 1% Tween® 20.

Ammonium in the buffer minimizes the need for optimization of the MgCl₂ concentration or the annealing temperature for most primer-template systems.

25 mM MgCl₂

Recommended Storage and Stability

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label.

Option: Store at +4 °C for up to 6 months.

Quality Control

Taq DNA Polymerase is tested for contaminating activities, with no traces of endonuclease activity, nicking activity or exonuclease activity.

Unit Definition

One unit is defined as the amount of polymerase that incorporates 10 nmol of dNTPs into acid-precipitable DNA in 30 minutes at 72 °C under standard assay conditions.

Protocol

This protocol serves as a guideline to ensure optimal PCR results when using Taq DNA Polymerase. Optimal reaction conditions such as incubation times, temperatures and amount of template DNA may vary and must be determined individually.

1. Thaw Solutions. **It is important to thaw all solutions completely (some buffers need to reach room temperature) and mix thoroughly before use to avoid localized concentrations of salts.** Keep all components on ice.
2. Set up reaction mixtures in an area separate from that used for DNA preparation or product analysis. Work on ice at all times.
3. Prepare a master mix according to Table 1. The master mix typically contains all the components needed for extension except the template DNA.

Table 1. Reaction components (master mix and template DNA)

Component	Vol./reaction*	Final concentration*
10x Buffer	5 μl	1x
25 mM MgCl ₂	0 μl (0 – 6 μl)	1.5 mM (1.5 – 4.5 mM)
dNTP mix (12.5 mM each)	0.8 μl	0.2 mM of each dNTP
Primer A (10 μM)	1 μl (0.5 – 5 μl)	0.2 μM (0.1 – 1.0 μM)
Primer B (10 μM)	1 μl (0.5 – 5 μl)	0.2 μM (0.1 – 1.0 μM)
Taq DNA Pol.	0.2 μl (0.2 – 1 μl)	1 unit (1 – 5 units)
PCR-grade H ₂ O	X μl	-
Template DNA	X μl	genomic DNA: 50 ng (10 – 500 ng) plasmid DNA: 0.5 ng (0.1 – 1 ng) bacterial DNA: 5 ng (1 – 10 ng)
TOTAL volume	50 μl	-

* Suggested starting conditions; theoretically used conditions in brackets. The final volume can be reduced to 25 μl by using half of the volumes suggested in Vol./reaction, eg. 0.1 μl Taq instead of 0.2 μl Taq.

4. Mix the master mix thoroughly and dispense appropriate volumes into reaction tubes. Mix gently, e.g. by pipetting the master mix up and down a few times.
5. Add template DNA to the individual tubes containing the master mix.
6. Program the thermal cycler according to the manufacturer's instructions.
For maximum yield and specificity, temperatures and cycling times should be optimized for each new template target or primer pair.
7. Place the tubes in the thermal cycler and start the reaction.

Three-step PCR program

Cycles	Duration of cycle	Temperature
1	2 – 5 minutes ^a	95 °C
25 – 35	20 – 30 seconds ^b 20 – 40 seconds ^c 30 seconds ^d	95 °C 50 – 65 °C 72 °C
1	5 minutes ^e	72 °C

^a. Initial denaturation step (optional).

^b. Denaturation step: This step is the first regular cycling event and consists of heating the reaction to 95 °C for 20 – 30 seconds. It causes melting of the DNA template by disrupting the hydrogen bonds between complementary bases, yielding single-stranded DNA molecules.

^c. Annealing step: The reaction temperature is lowered to 50 – 65 °C for 20 – 40 seconds allowing annealing of the primers to the single-

stranded DNA template. Typically, the annealing temperature is about 3 – 5 °C below the T_m (melting temperature) of the primers used.

- ^d Extension/elongation step: Taq polymerase has its optimum activity temperature at 72 °C. At this step the DNA polymerase synthesizes a new DNA strand complementary to the DNA template strand. The extension time depends on the length of the DNA fragment to be amplified. As a rule of thumb, at its optimum temperature the DNA polymerase will polymerize a thousand bases per minute.
- ^e Final elongation: This single step is occasionally performed at a temperature of 72 °C for 5 minutes after the last PCR cycle to ensure that any remaining single-stranded DNA is fully extended.

Two-step PCR program

Fast 2-step PCR protocols are available using this link:

<https://ampliqon.com/en/pcr-technology/application-notes/>

Notes:

- 15 mM $MgCl_2$ is present in 10x PCR Buffer. The 1x concentration is 1.5 mM $MgCl_2$. In some applications, more than 1.5 mM $MgCl_2$ is required for best results. For this reason, 25 mM $MgCl_2$ is included in the kit. Table 2 provides the volume of 25 mM $MgCl_2$ to be added to the master mix if a higher $MgCl_2$ concentration is required.

Table 2. Additional volume (μl) of $MgCl_2$ per 50 μl reaction

Final $MgCl_2$ conc. in reaction (mM)	1.5	2.0	2.5	3.0	3.5	4.0	4.5
Volume of 25 mM $MgCl_2$	0	1	2	3	4	5	6

- For longer DNA targets more DNA polymerase could be added to the PCR master mix.

Related Products

Taq DNA Polymerase (500 units) *	Cat. No.
Taq DNA Polymerase 5 U/μl	A110003
• with 10x Ammonium Buffer	A111103
• 5x PCR Buffer RED	A111803
Taq DNA Polymerase 5 U/μl, RED	A200003
• with 10x Ammonium Buffer	A201103
Taq DNA Polymerase 5 U/μl, glycerol free	A100003
• with 10x Ammonium Buffer	A101103

Hot Start DNA Polymerase (500 units) *	Cat. No.
TEMPase Hot Start DNA Polymerase, 5 U/μl	A220003
• with 10x Ammonium Buffer	A221103

*Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM $MgCl_2$.

Buffers for DNA polymerases *	Cat. No.
10x Ammonium Buffer, 3 x 1.5 ml	A301103
10x Standard Buffer, 3 x 1.5 ml	A302103
10x Combination Buffer, 3 x 1.5 ml	A303103
5x PCR Buffer RED, 6 x 1.5 ml **	A301810

*Ammonium Buffer, Standard Buffer and Combination Buffer are also available as Mg^{2+} free buffers, detergent free buffers and Mg^{2+} and detergent free buffers.

**For direct gel loading and visualisation.

Taq 2x Master Mixes (500 x 50 μl reactions) *	Cat. No.
Taq OptiMix Mix Clear 2x Master Mix	A370503
Taq DNA Polymerase 2x Master Mix	A140303
Taq DNA Polymerase 2x Master Mix RED	A180303

*Taq Master Mixes available also in 1.1x variants.

Ultrapure dNTPs*	Cat. No.
dNTP Mix 40 mM (2 x 500 μl): 10 mM each dA, dC, dG, dT	A502004
dNTP Set, 100 mM each: 250 μl of each dA, dC, dG and dT	A511104

*Other concentrations and Single dNTPs are available.

Loading Buffers and Ladders	Cat. No.
5x Loading Buffer Red *, 5 x 1 ml	A608104
Iqon PCR Ladder **, 100 – 3000 bp, 1 x 0.5 ml	A610341

* Also available with Blue, Orange or Cyan. ** Available in different size ranges.

For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at www.ampliqon.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.

Made in Denmark

Issued 08/2021