

Taq DNA Polymerase 2x Master Mix

2 mM MgCl₂ final concentration

Cat. No.: A150301



100 Reactions

MADE IN DENMARK

| | |
|------------|---|
| - | Taq DNA Polymerase 2x Master Mix, 2 mM MgCl ₂ |
| ID No. | 5200150 |
| Cap colour | Purple |
| Content | 2 x 1.25 ml |

Key Features

Taq DNA Polymerase 2x Master Mix is a ready-to-use 2x reaction mix with the Ampliqon Taq DNA polymerase, the NH₄⁺ buffer system, dNTPs and magnesium chloride present. Each reaction requires 25 µl of the 2x Master Mix. Simply add primers, template and water to a total reaction volume of 50 µl to successfully carry out primer extensions and other molecular biology applications.

Taq DNA Polymerase 2x Master Mix offers several advantages. Set up time is significantly reduced. The chance of contaminating component stocks is eliminated. Reduction of reagent handling steps leads to better reproducibility. Standard tests can be set up with the confidence that results will be consistent every time.

Composition of the Taq DNA Polymerase 2x Master Mix (2 mM MgCl₂ final concentration)

- Tris-HCl pH 8.5, (NH₄)₂SO₄, 4 mM MgCl₂, 0.2% Tween® 20
- 0.4 mM of each dNTP
- Ampliqon Taq DNA polymerase
- Stabilizer

Recommended Storage and Stability

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label.

Option: Store at +4 °C for up to 6 months.

Quality Control

Taq DNA Polymerase is tested for contaminating activities, with no traces of endonuclease activity, nicking activity or exonuclease activity.

Protocol

This protocol serves as a guideline to ensure optimal PCR results when using Taq DNA Polymerase 2x Master Mix. Optimal reaction conditions such as incubation times, temperatures, and amount of template DNA may vary and must be determined individually.

1. Thaw Taq 2x Master Mix and primers. **It is important to thaw the solutions completely and mix thoroughly before use to avoid localized concentrations of salts.** Keep all components on ice.
2. Set up each reaction. Table 1 shows the reaction set up for a final volume of 50 µl. If desired, the reaction size may be scaled down. Use 10 µl of the Taq 2x Master Mix in a final volume of 20 µl.

Table 1. Reaction components (reaction mix and template DNA)

| Component | Vol./reaction* | Final concentration* |
|----------------------------|-------------------|---|
| Taq 2x Master Mix | 25 µl | 1x |
| 25 mM MgCl ₂ | 0 µl (0 – 5 µl) | 2 mM (2 – 4.5 mM) |
| Primer A (10 µM) | 1 µl (0.5 – 5 µl) | 0.2 µM (0.1 – 1.0 µM) |
| Primer B (10 µM) | 1 µl (0.5 – 5 µl) | 0.2 µM (0.1 – 1.0 µM) |
| PCR-grade H ₂ O | X µl | - |
| Template DNA | X µl | genomic DNA: 50 ng (10 – 500 ng) plasmid DNA: 0.5 ng (0.1 – 1 ng) bacterial DNA: 5 ng (1 – 10 ng) |
| TOTAL volume | 50 µl | - |

* Suggested starting conditions; theoretically used conditions in brackets

3. Mix the reaction mix thoroughly and dispense appropriate volumes into reaction tubes. Mix gently, e.g. by pipetting the reaction mix up and down a few times.
4. Add template DNA to the individual tubes containing the reaction mix.
5. Program the thermal cycler according to the manufacturer's instructions. See table 2 for an example.
For maximum yield and specificity, temperatures and cycling times should be optimized for each new template target or primer pair.
6. Place the tubes in the thermal cycler and start the reaction.

Table 2. Three-step PCR program

| Cycles | Duration of cycle | Temperature |
|---------|---|------------------------------|
| 1 | 2 – 5 minutes | 95 °C |
| 25 - 35 | 20 – 30 seconds ^a 20 – 40 seconds ^b 30 seconds ^c | 95 °C 50 – 65 °C 72 °C |
| 1 | 5 minutes ^d | 72 °C |

^a. Denaturation step: This step is the first regular cycling event and consists of heating the reaction to 95 °C for 20 – 30 seconds. It causes melting of the DNA template by disrupting the hydrogen bonds between complementary bases, yielding single-stranded DNA molecules.

^b. Annealing step: The reaction temperature is lowered to 50 – 65 °C for 20 – 40 seconds allowing annealing of the primers to the single-

stranded DNA template. Typically, the annealing temperature is about 3 – 5 °C below the T_m (melting temperature) of the primers used.

- ^c Extension/elongation step: Taq polymerase has its optimum activity temperature at 72 °C. At this step the DNA polymerase synthesizes a new DNA strand complementary to the DNA template strand. The extension time depends on the length of the DNA fragment to be amplified. As a rule of thumb, at its optimum temperature the DNA polymerase will polymerize a thousand bases per minute.
- ^d Final elongation: This single step is occasionally performed at a temperature of 72 °C for 5 minutes after the last PCR cycle to ensure that any remaining single-stranded DNA is fully extended.

Two-step PCR program

Fast 2-step PCR protocols are available using this link:
<https://ampliqon.com/en/pcr-technology/application-notes/>

Notes:

- The final $MgCl_2$ concentration of this 2x Taq Master Mix is 2 mM. In some applications, more than 2 mM $MgCl_2$ is required for best results. Use 25 mM $MgCl_2$ to adjust the Mg^{2+} concentration according to table 3.

Table 3. Additional volume (μ l) of $MgCl_2$ per 50 μ l reaction:

| Final $MgCl_2$ conc. in reaction (mM) | 2.0 | 2.5 | 3.0 | 3.5 | 4.0 | 4.5 |
|---------------------------------------|-----|-----|-----|-----|-----|-----|
| Volume of 25 mM $MgCl_2$ | 0 | 1 | 2 | 3 | 4 | 5 |

Related Products

| Taq Master Mixes (500 x 50 μ l reactions) * | Cat. No. |
|---|----------------|
| 2x Master Mix, 1.5 mM $MgCl_2$ final concentration | A140303 |
| 2x Taq OptiMix CLEAR, 1.5 mM $MgCl_2$ final concentration | A370503 |
| 2x Master Mix RED, 1.5 mM $MgCl_2$ final concentration | A180303 |

| TEMPase Hot Start Master Mixes (500 x 50 μ l reactions) * | Cat. No. |
|---|----------------|
| 2x Master Mix A**, 1.5 mM $MgCl_2$ final concentration | A230303 |
| 2x Master Mix A**BLUE, 1.5 mM $MgCl_2$ final concentration | A290403 |

*Master mixes available also in 1.1x variants as well as 2 mM $MgCl_2$ variants, **Mix A is Ammonium Buffer based, also available as Mix C based on Combination Buffer.

| Special TEMPase Master Mixes (500 x 50 μ l reactions) | Cat. No. |
|--|----------------|
| Multiplex 2x Master Mix, 3 mM $MgCl_2$ final concentration | A260303 |
| GC TEMPase 2x Master Mix I – for GC-rich templates | A331703 |
| GC TEMPase 2x Master Mix II – for GC-rich templates | A332703 |

| Taq DNA Polymerase (500 units) * | Cat. No. |
|----------------------------------|----------------|
| Taq DNA Polymerase 5 U/ μ l | A110003 |
| ● with 10x Ammonium Buffer | A111103 |

*Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM $MgCl_2$

| Hot Start DNA Polymerase (500 units) * | Cat. No. |
|--|----------------|
| TEMPase Hot Start DNA Polymerase, 5 U/ μ l | A220003 |
| ● with 10x Ammonium Buffer | A221103 |

*Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM $MgCl_2$

| Buffers for DNA polymerases * | Cat. No. |
|------------------------------------|----------------|
| 10x Ammonium Buffer, 3 x 1.5 ml | A301103 |
| 10x Standard Buffer, 3 x 1.5 ml | A302103 |
| 10x Combination Buffer, 3 x 1.5 ml | A303103 |
| 5x PCR Buffer RED, 6 x 1,5 ml ** | A301810 |
| PCR Grade Water, 6 x 5 ml | A360056 |

*Ammonium Buffer, Standard Buffer and Combination Buffer are also available as Mg^{2+} free buffers, detergent free buffers and Mg^{2+} and detergent free buffers.

**For direct gel loading and visualisation.

Reagents for *in vitro* laboratory use only.

Other product sizes, combinations and customized solutions are available. Please look at www.ampliqon.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.

Made in Denmark

Issued 10/2019