

# Standard Buffer (Mg<sup>2+</sup> free)

Cat. No.: A302210

MADE IN **DENMARK** 

-	10x Standard Buffer, Mg <sup>2+</sup> free
ID No.	5100620
Cap color	White
Content	10 x 1.5 ml

#### **Features and General Description**

10x Standard Buffer are usually supplied in 10x formulations with 15 mM  $MgCl_2$  included but are also available as  $Mg^{2+}$  free buffer, detergent free buffer as well as  $Mg^{2+}$  and detergent free buffer.

#### **Standard Buffer**

Standard Buffer is the traditional potassium ( $K^+$ ) buffer. Standard Buffer promotes high specificity and careful optimization of primer annealing temperatures and  $Mg^{2+}$  concentrations may be required.

#### Magnesium

Mg<sup>2+</sup> is required for polymerase activity. Low Mg<sup>2+</sup> concentrations increase the fidelity but with too low Mg<sup>2+</sup> concentrations the polymerase will not work. The Mg<sup>2+</sup> concentration available in the reaction is dependent on several parameters e.g. the presence of chelators or the dNTP concentration. Therefore, the Mg<sup>2+</sup> concentration should be optimized.

#### Tween

Non-ionic detergents are used to prevent the polymerase to stick to the walls of the tube, to stabilize the polymerase and increase yield. However, these agents might increase non-specific amplification or interfere with downstream reactions. Tween can be used to neutralize SDS contaminations in the DNA template.

### **Recommended Storage and Stability**

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label

Option: Store at +4 °C for up to 6 months.

## **Quality Control**

Each lot of buffer is functionally tested in PCR.

# **Kit Components**

10x Standard Buffer, Mg<sup>2+</sup> free
Tris-HCl pH 8.5, KCl, 1% Tween<sup>®</sup> 20.

# Determining the optimal buffer system for your application

Ampliqon offers several PCR buffers to allow the customer to choose the optimal buffer system for a specific amplification process.

For your specific application the optimal reaction condition can be determined by comparing PCR reactions containing the different Ampliqon buffers.

The final concentration of the buffer in the reaction should be 1x.




Figure 1: Amplification of three different cDNA templates using Ammonium Buffer versus Standard Buffer.

#### **Related Products**

Taq DNA Polymerase (500 units) *	Cat. No.
Taq DNA Polymerase 5 U/μl	A110003
• with 10x Ammonium Buffer	A111103
• 5x PCR Buffer RED	A111803
Tag DNA Polymerase 5 U/μl, glycerol free	A100003
• with 10x Ammonium Buffer	A101103
Had Clark DNA Dalamana (FOO and Lab #	0.1.11.
Hot Start DNA Polymerase (500 units) *	Cat. No.

Hot Start DNA Polymerase (500 units) *	Cat. No.
TEMPase Hot Start DNA Polymerase, 5 U/μl	A220003
with 10x Ammonium Buffer	A221103

<sup>\*</sup>Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM  $\rm MgCl_2$ .

Cat. No.
A301103
A302103
A303103
A301810
A301703
A302703

<sup>\*</sup>Ammonium Buffer, Standard Buffer and Combination Buffer are also available as Mg<sup>2+</sup> free buffers, detergent free buffers and Mg<sup>2+</sup> and detergent free buffers.

\*\*For direct gel loading and visualisation.

Ultrapure dNTPs*	Cat. No.
dNTP Mix 40 mM (2 x 500 μl): 10 mM each dA, dC, dG, dT	A502004
dNTP Set, 100 mM each: 250 μl of each dA, dC, dG and dT	A511104

<sup>\*</sup>Other concentrations and Single dNTPs are available.

Loading Buffers, PCR Water and Ladders	Cat. No.
5x Loading Buffer Red *, 5 x 1 ml	A608104
Iqon PCR Ladder **, 100 – 3000 bp, 1 x 0.5 ml	A610341
PCR Grade Water, 6 x 5 ml	A360056

<sup>\*</sup> Also available with Blue, Orange or Cyan. \*\* Available in different size ranges. For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at www.ampliqon.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.