# AMPLIQON III

### Standard Buffer (Mg<sup>2+</sup> free, Tween free)

Cat. No.: A302556

## A302556

ENMAR	-	10x Standard Buffer, Mg <sup>2+</sup> free, Tween free
IN D	ID No.	5100650
)E II	Cap color	Clear
MAI	Content	6 x 5 ml

#### **Features and General Description**

10x Standard Buffer are usually supplied in 10x formulations with 15 mM  $MgCl_2$  included but are also available as  $Mg^{2^+}$  free buffer, detergent free buffer as well as  $Mg^{2^+}$  and detergent free buffer.

#### **Standard Buffer**

Standard Buffer is the traditional potassium ( $K^+$ ) buffer. Standard Buffer promotes high specificity and careful optimization of primer annealing temperatures and Mg<sup>2+</sup> concentrations may be required.

#### Magnesium

 $Mg^{2+}$  is required for polymerase activity. Low  $Mg^{2+}$  concentrations increase the fidelity but with too low  $Mg^{2+}$  concentrations the polymerase will not work. The  $Mg^{2+}$  concentration available in the reaction is dependent on several parameters e.g. the presence of chelators or the dNTP concentration. Therefore, the  $Mg^{2+}$  concentration should be optimized.

#### Tween

Non-ionic detergents are used to prevent the polymerase to stick to the walls of the tube, to stabilize the polymerase and increase yield. However, these agents might increase nonspecific amplification or interfere with downstream reactions. Tween can be used to neutralize SDS contaminations in the DNA template.

#### **Recommended Storage and Stability**

Long term storage at -20 °C. Product expiry at -20 °C is stated on the label.

Option: Store at +4 °C for up to 6 months.

Quality Control Each lot of buffer is functionally tested in PCR.

#### **Kit Components**

**10x Standard Buffer, Mg<sup>2+</sup> free, Tween free** Tris-HCl pH 8.5, KCl.

### Determining the optimal buffer system for your application

Ampliqon offers several PCR buffers to allow the customer to choose the optimal buffer system for a specific amplification process.

For your specific application the optimal reaction condition can be determined by comparing PCR reactions containing the different Ampliqon buffers.

The final concentration of the buffer in the reaction should be 1x.




Figure 1: Amplification of three different cDNA templates using Ammonium Buffer versus Standard Buffer.

#### **Related Products**

Taq DNA Polymerase (500 units) *	Cat. No.	
Taq DNA Polymerase 5 U/μl	A110003	
<ul> <li>with 10x Ammonium Buffer</li> </ul>	A111103	
• 5x PCR Buffer RED	A111803	
Taq DNA Polymerase 5 U/μl, glycerol free	A100003	
with 10x Ammonium Buffer	A101103	
Hot Start DNA Polymerase (500 units) *	Cat. No.	
TEMPase Hot Start DNA Polymerase, 5 U/μl	A220003	
<ul> <li>with 10x Ammonium Buffer</li> </ul>	A221103	
*Available in kits including one or two buffers (Ammonium Puffer, Standard Puffer		

\*Available in kits including one or two buffers (Ammonium Buffer, Standard Buffer or Combination Buffer). All kits include extra 25 mM MgCl<sub>2</sub>.

Buffers for DNA Polymerases *	Cat. No.
10x Ammonium Buffer, 3 x 1.5 ml	A301103
10x Standard Buffer, 3 x 1.5 ml	A302103
10x Combination Buffer, 3 x 1.5 ml	A303103
5x PCR Buffer RED, 6 x 1,5 ml **	A301810
4x GC Buffer I, 3 x 1.5 ml	A301703
4x GC Buffer II, 3 x 1.5 ml	A302703

\*Ammonium Buffer, Standard Buffer and Combination Buffer are also available as  $Mg^{2^+}$  free buffers, detergent free buffers and  $Mg^{2^+}$  and detergent free buffers. \*\*For direct gel loading and visualisation.

Ultrapure dNTPs*	Cat. No.
dNTP Mix 40 mM (2 x 500 μl): 10 mM each dA, dC, dG, dT	A502004
dNTP Set, 100 mM each: 250 $\mu l$ of each dA, dC, dG and dT	A511104
*Other concentrations and Single dNTPs are available.	

Loading Buffers, PCR Water and Ladders	Cat. No.
5x Loading Buffer Red *, 5 x 1 ml	A608104
Iqon PCR Ladder **, 100 – 3000 bp, 1 x 0.5 ml	A610341
PCR Grade Water, 6 x 5 ml	A360056

\* Also available with Blue, Orange or Cyan. \*\* Available in different size ranges. For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at www.ampliqon.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.