

AQ90 High Fidelity DNA Polymerase 2x Master Mix

MADE IN DENMARK

Cat. No.: A470701
100 Reactions



R470701

| AQ90 High Fidelity DNA Polymerase 2x Master Mix 4 mM MgCl ₂ | |
|---|-------------|
| ID No. | 5500400 |
| Cap colour | Purple |
| Content | 2 x 1.25 ml |

Key Features

- Convenient reaction set-up
- High fidelity – measured up to 50x *Taq* fidelity
- Excellent coverage for amplification of difficult amplicons with low to high GC content
- Long range capability: 8.5 kb for gDNA and ≤ 12.5 kb for λDNA
- Recommended for cloning, mutagenesis and other molecular applications requiring extremely high fidelity

AQ90 High Fidelity DNA Polymerase 2x Master Mix is an all-in-one 2x master mix containing the AQ90 High Fidelity DNA Polymerase, AQ90 Buffer, dNTPs and MgCl₂. Simply mix AQ90 High Fidelity DNA Polymerase 2x Master Mix with primers, DNA template and water and you are ready to carry out PCR.

AQ90 High Fidelity DNA Polymerase is a thermostable DNA Polymerase with proofreading ability. AQ90 High Fidelity DNA Polymerase exhibits 5'→3' DNA polymerase activity and 3'→5' proofreading exonuclease activity. The latter allows the enzyme to correct misincorporated nucleotides. The AQ90 High Fidelity DNA Polymerase 2x Master Mix exhibits robust amplification of targets with low to high GC content, as well as a fidelity* up to 50x *Taq*.

*Estimated by NGS technology using the Illumina MiSeq instrument.

Protocol

Reaction conditions such as incubation times, temperatures and amount of template DNA may vary and must be determined individually. Amplification of templates with high GC content, high secondary structures as well as long range amplification may require more optimization.

Set up reaction mixtures in an area separate from that used for DNA preparation or product analysis. **Work on ice at all times.**

1. Thaw AQ90 High Fidelity DNA Polymerase 2x Master Mix and primer solutions.
It is recommended to completely thaw and thoroughly mix the master mix to ensure proper resuspension of precipitates.

2. Prepare the reaction mix. Table 1 shows the reaction set up for a final volume of 50 µl. If desired, the reaction size may be scaled down.

Table 1. Reaction components

| Component | Vol./reaction* | Final concentration* |
|----------------------------|-----------------|---|
| 2x Master Mix | 25 µl | 1x |
| Primer A (10 µM) | 1 µl | 0.2 µM |
| Primer B (10 µM) | 1 µl | 0.2 µM |
| 25 mM MgCl ₂ | 0 µl (0 – 6 µl) | 2 mM (2 – 5 mM) |
| Betaine (5M) (optional) | 10 – 20 µl | 1 – 2M |
| PCR-grade H ₂ O | X µl | - |
| Template DNA | X µl | genomic DNA: 50 ng (10 – 500 ng) plasmid DNA: 0.5 ng (0.1 – 1 ng) bacterial DNA: 5 ng (1 – 10 ng) |
| TOTAL volume | 50 µl | - |

* Suggested starting conditions; theoretically used conditions in brackets.

3. Mix the reaction mix thoroughly and dispense appropriate volumes into reaction tubes. Mix gently, e.g. by pipetting the master mix up and down a few times.
4. Add template DNA to the individual tubes containing the reaction mix.
5. Program the thermal cycler according to the manufacturer's instructions. For maximum yield and specificity, temperatures and cycling times should be optimized for each new template target or primer pair.
6. Place the tubes in the thermal cycler and start the reaction.

Table 2. Three-step PCR program

| Cycles | Duration of cycle | Temperature |
|---------|---|------------------------------|
| 1 | 2 – 5 minutes ^a | 95 °C |
| 25 – 35 | 10 – 30 seconds ^b 20 – 40 seconds ^c 30 seconds ^d | 95 °C 55 – 70 °C 72 °C |
| 1 | 5 minutes ^e | 72 °C |

^a. Initial denaturation step (optional).

^b. Denaturation step: This step is the first regular cycling event and consists of heating the reaction to 95 °C for 10 – 30 seconds. It causes melting of the DNA template by disrupting the hydrogen bonds between complementary bases, yielding single-stranded DNA molecules.

^c. Annealing step: The reaction temperature is lowered to 55 – 70 °C for 20 – 40 seconds allowing annealing of the primers to the single-stranded DNA template. Typically, the annealing temperature is about 3 – 5 °C below the T_m (melting temperature) of the primers used. **Because of the high salt content within the AQ90 High Fidelity DNA Polymerase 2x Master Mix, annealing temperature will likely be higher than with more traditional PCR master mixes.**

^d. Extension/elongation step: AQ90 High Fidelity DNA polymerase has its optimum activity temperature at 72 °C. At this step the DNA polymerase synthesizes a new DNA strand complementary to the DNA template strand. The extension time depends on the length of the DNA fragment to be amplified. **Generally, we recommend an extension time of 1 minute per kb, especially for longer amplicons.**

^e. Final elongation: This single step is occasionally performed at a temperature of 72 °C for 2 – 5 minutes after the last PCR cycle to ensure that any remaining single-stranded DNA is fully extended.

Notes for optimization of PCR conditions:

- The optimal MgCl_2 concentration should be determined empirically, but in most cases a final concentration of 2 mM, as provided in AQ90 High Fidelity DNA Polymerase 2x Master Mix, will produce satisfactory results. Table 3 provides the volume of 25 mM MgCl_2 to be added to the master mix if a higher MgCl_2 concentration is required.

Table 3. Additional volume (μl) of MgCl_2 per 50 μl reaction

| Final MgCl_2 conc. in reaction (mM) | 2.0 | 2.5 | 3.0 | 3.5 | 4.0 | 4.5 | 5 |
|--|-----|-----|-----|-----|-----|-----|---|
| Volume of 25 mM MgCl_2 | 0 | 1 | 2 | 3 | 4 | 5 | 6 |

- For difficult amplicons, such as GC-rich templates, those with secondary structures or very long amplicons the addition of 1 - 2 M Betaine solution might improve reaction performance (See Additional Products for ordering information). 5 - 10 % DMSO can also be used to improve reaction performance. If using high concentrations of DMSO the annealing temperature has to be lowered as it decreases the primer T_m .
- Primers of 20 – 40 nucleotides with a GC content of 40 - 60 % are recommended. Online Software such as the Primer3plus <https://primer3plus.com/cgi-bin/dev/primer3plus.cgi> can be used to design primers.

Kit Components

AQ90 High Fidelity DNA Polymerase 2x Master Mix

- AQ90 High Fidelity DNA Polymerase
- Optimized buffer components, 4.0 mM MgCl_2
- dNTPs

5 M Betaine Enhancer Solution

Sold separately. Cat No.: A351104

Recommended Storage and Stability

Long term storage at $-20\text{ }^{\circ}\text{C}$. Product expiry at $-20\text{ }^{\circ}\text{C}$ is stated on the label.

Optional: Store at $+4\text{ }^{\circ}\text{C}$ for up to 6 months.

Quality Control

AQ90 High Fidelity DNA Polymerase is tested for contaminating activities with no traces of endonuclease activity or nicking activity.

Unit Definition

One unit is defined as the amount of enzyme that incorporates 10 nmoles of dNTPs into an acid-precipitable form of DNA in 30 minutes at $72\text{ }^{\circ}\text{C}$ under standard assay conditions.

Related Products

| AQ90 High Fidelity DNA Polymerase 2x Master Mix | Cat. No. |
|---|----------|
| 100 reactions | A470701 |
| 500 reactions | A470703 |
| 2500 reactions | A470706 |
| 5000 reactions | A470707 |

| AQ90 High Fidelity DNA Polymerase 2 U/ μl | Cat. No. |
|--|----------|
| With 10x AQ90 Buffer – 100 U | A457401 |
| With 10x AQ90 Buffer – 500 U | A457403 |
| With 10x AQ90 Buffer – 1000 U | A457404 |
| With 10x AQ90 Buffer – 2500 U | A457406 |

| PCR Grade Water | Cat. No. |
|-----------------|----------|
| 6 x 5 ml | A360056 |

| Betaine Enhancer Solution 5 M | Cat. No. |
|-------------------------------|----------|
| 5 x 1 ml | A351104 |

For Research Use Only. Not for use in diagnostics procedures.

Other product sizes, combinations and customized solutions are available. Please look at www.ampliqon.com or ask for our complete product list for PCR Enzymes. For customized solutions please contact us.

Made in Denmark

Issued 08/2021